

RESEARCH ARTICLE

Open Access

No association between polymorphisms of *WNT2* and schizophrenia in a Korean population

Hak-Jae Kim¹, Jin Kyung Park², Su Kang Kim³, Sung Wook Kang³, Jong Woo Kim², Hyun-Kyung Park⁴, Ah-Rang Cho², Ji Young Song² and Joo-Ho Chung^{*3}

Abstract

Background: Wingless-type MMTV integration site family member 2 (*WNT2*) has a potentially important role in neuronal development; however, there has yet to be an investigation into the association between single nucleotide polymorphisms (SNPs) of *WNT2* and schizophrenia. This study aimed to determine whether certain SNPs of *WNT2* were associated with schizophrenia in a Korean population.

Methods: We genotyped 7 selected SNPs in the *WNT2* gene region (approximately 46 Kb) using direct sequencing in 288 patients with schizophrenia and 305 healthy controls.

Results: Of the SNPs examined, one SNP showed a weak association with schizophrenia ($p = 0.017$ in the recessive model). However, this association did not remain statistically significant after Bonferroni correction.

Conclusion: The present study does not support a major role for *WNT2* in schizophrenia. This could be due to the size of the population. Therefore, additional studies would be needed to definitively rule out the gene's minor effects.

Background

Schizophrenia is a complex disease that develops in approximately 0.5-1.0% of the global population and inflicts physical suffering and mental anguish on the affected individuals and their families [1,2]. The neurodevelopmental theory of schizophrenia postulates that abnormal neurodevelopment is one of the etiological factors of schizophrenia [3-6]. Recent studies suggest that schizophrenia may be a subtle disorder of brain development or plasticity [7-9].

Wnt proteins belong to a family of molecules that locally activate cell signaling pathways, regulating cell fate and playing an important role in development [10]. Evidence suggests that Wnt signaling and insulin signaling, which regulates glycogen synthase kinase-3 (GSK-3), may be important in schizophrenia [11]. Several lines of evidence support the association between signaling aberrations and schizophrenia. However, research has shown changes in gene expressions in schizophrenia [12-16], and, in particular, that some such changes affect GSK-3 [14,17-19]. Researchers have also found Wnt signaling

abnormalities associated with schizophrenia. For example, expressions of β -catenin proteins decreased in the brains of schizophrenia patients, presumably due to aberrant regulation of β -catenin degradation mediated by activated GSK-3 β [20,21]. Another report suggested the enhanced GSK-3 β activity was secondary to diminished phosphorylation by Akt, since there is decreased Akt expression in schizophrenia patients' brains [22]. Furthermore, researchers found increased Wnt-1 expression in the hippocampi of schizophrenia patients relative to normal subjects. Wnt-1 is an important molecule in the Wnt pathway [23]. Several studies have implicated Wnt signaling genes in bipolar disorder, as well as in schizophrenia. Researcher found that the frizzled-3 gene (*FZD3*), which encodes a receptor for Wnt ligands, was associated with schizophrenia in 3 different samples [24-26].

The wingless-type MMTV integration site family, member 2 (*WNT2*) gene is located at chromosome 7q31.2, a region that, in a genome scan study, showed evidence of a link to schizophrenia [27]. *WNT2* is one of the *WNT* genes that are expressed in a variety of tissues during development [28]. Knockout and expression studies in mice, zebra fish, and *Xenopus* have demonstrated *WNT* genes' crucial roles in the development and pat-

* Correspondence: jhchung@khu.ac.kr

³ Kohwang Medical Research Institute, School of Medicine, Kyung Hee University, Seoul 130-701, Republic of Korea

Full list of author information is available at the end of the article

tering of the central nervous system [29-31]. WNT genes encode several signaling proteins that are essential in embryo patterning, cell proliferation, and cell determination [32,33]. The WNT signaling pathway also includes several receptors expressed in both developing and mature nervous systems [34].

Previously, several studies have addressed whether the *WNT2* gene is associated with autism, a neurodevelopmental disorder [35-37]. Although Wassink et al. reported that one of *WNT2* SNPs (rs2024233 on exon 5) is associated with autism [35], following association studies did not find any significant associations with autism [36,37]. In addition, Proitsi et al. performed a combined positional and candidate association screen, by identifying known *WNT* signaling genes, and showed that Dickkopf-4 was associated with schizophrenia [38]. There have been no published investigations of the association between *WNT2* genetic polymorphisms and schizophrenia, despite its potentially important developmental roles and its importance in genetics. In this study, therefore, we investigated the association between *WNT2* polymorphisms and schizophrenia in a Korean population.

Methods

Subjects

For this study, we evaluated 288 schizophrenic patients (193 males, aged 42.98 ± 10.63 years [mean \pm SD]; 95 females, aged 43.03 ± 10.65) and 305 controls with no clinical evidence of any disorders (147 males, 39.92 ± 5.82 ; 158 females, 36.55 ± 6.77). The schizophrenic patients met the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) [39] criteria for schizophrenia. We reviewed each patient's medical record and rated the schizophrenic patients using the Brief Psychiatric Rating Scale (BPRS) [40], the Scale for the Assessment of Negative Symptoms (SANS) [41], and the operational criteria (OPCRIT) checklist [42].

To evaluate patients for schizophrenia with poor concentration, we used the OPCRIT Checklist. Of the many OPCRIT items, we selected only a few items, already described in our previous study, for use [43]. We determined which patients to place in the poor concentration subgroup according to patients' subjective complaints of being unable to think clearly or make decisions (scoring: 1 for a duration of one week, 2 for two weeks, and 3 for one month) [42]. Thus, we examined 167 schizophrenic patients having poor concentration (116 males, 42.8 ± 10.0 ; 51 females, 41.9 ± 10.8) and 121 schizophrenic patients who had no such complaints (77 males, 43.6 ± 11.2 ; 44 females, 44.8 ± 10.7). We also analyzed the associations between the schizophrenic patients' SNPs and their SANS scores (Table 1). The genotype frequencies of rs6948009 was evaluated (A/A, $n = 211$; A/G, $n = 71$; G/G, $n = 3$) for the analysis with SANS scores.

We recruited control subjects who had been found mentally fit by a general health checkup program. All studies were carried out according to the guidelines of the Declaration of Helsinki [44]. We obtained written informed consent from each subject. The study was approved by the ethics review committee of the Medical Research Institute, Kyung Hee University Medical Center, Seoul, Republic of Korea.

SNP Genotyping

We selected 8 SNPs from *WNT2* genes in chromosome 7q31.2 by downloading all the SNPs typed in the *WNT2* genes from the HapMap database (<http://www.hapmap.org/>; genome build 35) and dbSNP database version 129. From these, we selected tag SNPs using the Tagger program's aggressive tagging option (Paul de Bakker, <http://www.broad.mit.edu/mpg/tagger/>), so as to capture SNPs with a minor allele frequency of $< 5\%$ ($r^2 > 0.8$). This gave 2 exonic SNPs (rs2024233 and rs1051751), 3 intronic SNPs (rs733154, rs3779548, and rs17132543), and 3 regulatory SNPs (thought to be the promoter; the intervals are 2000 bp up- and downstream between the genes) (rs6948009 [869 bp downstream from the last nucleotide of the 3'-UTR], rs4730775 [164 bp downstream from the last nucleotide of the 3'-UTR], and rs39315 [219 bp upstream from the first nucleotide of the 5'-UTR]). In addition, we chose these SNPs because they were previously evaluated for relation to schizophrenia risk or showed evidence of functional significance. Figure 1 shows the locations of the selected SNPs.

Using a commercially available Qiagen DNA Extraction kit (Qiagen, Tokyo, Japan), we extracted genomic DNA from blood samples that had been placed in EDTA. We amplified the genomic DNA using the primers shown in Table 2 for each SNP, sequenced the PCR products using an ABI Prism 377 automatic sequencer (PE Applied Biosystems, Foster City, California, USA), and analyzed the sequence data using SeqManII software (DNASTAR Inc., Madison, Wisconsin, USA).

Statistical analysis

Using SNPstats, we assessed the Hardy-Weinberg equilibrium (HWE) for both controls and schizophrenia patients [45]. For the linkage disequilibrium (LD) block, we used the Haploview version 3.32 [46]. We inferred the haplotypes and their frequencies using the EM algorithm [47]. We used the multiple logistic regression model to calculate odds ratios (OR), 95% confidence intervals, and corresponding p values (controlling age and gender as covariables), to analyze the association between schizophrenia and both SNPs and haplotypes. Furthermore, we analyzed the association between the schizophrenia subgroup and both SNPs and haplotypes, using SNPstats, the HapAnalyzer version 1.0 [48], and HelixTree (Golden

Table 1: WNT2 SNPs genotype and allele frequencies in schizophrenia patients and healthy controls

SNP	Genotype	Schizophrenia		Control		Model	OR (95% CI)	P value
		Freq	%	Freq	%			
rs39315 5'-near gene	A/A	90	0.31	79	0.26	co-dominant	1.21 (0.95-1.54)	0.12
	A/G	137	0.48	150	0.49	dominant	1.38 (0.94-2.02)	0.1
	G/G	61	0.21	76	0.25	recessive	1.20 (0.80-1.81)	0.38
rs17132543 Intron4	A/A	97	0.34	98	0.32	co-dominant	1.14 (0.89-1.45)	0.3
	A/G	138	0.48	145	0.48	dominant	1.16 (0.80-1.68)	0.43
	G/G	53	0.18	62	0.2	recessive	1.22 (0.79-1.89)	0.37
rs3779548 Intron4	A/A	127	0.44	135	0.44	co-dominant	0.97 (0.74-1.25)	0.8
	A/G	128	0.44	136	0.45	dominant	0.99 (0.70-1.40)	0.96
	G/G	33	0.11	34	0.11	recessive	0.88 (0.50-1.53)	0.65
rs733154 Intron4	A/A	171	0.59	177	0.58	co-dominant	1.00 (0.74-1.33)	0.97
	A/G	103	0.36	109	0.36	dominant	1.02 (0.72-1.44)	0.93
	G/G	14	0.05	19	0.06	recessive	0.89 (0.42-1.92)	0.78
rs2024233 3'-UTR	A/A	87	0.3	84	0.28	co-dominant	1.11 (0.87-1.43)	0.41
	A/G	148	0.51	157	0.51	dominant	1.15 (0.79-1.69)	0.46
	G/G	53	0.18	64	0.21	recessive	1.14 (0.74-1.77)	0.54
rs4730775 3'-near gene	G/G	176	0.61	184	0.6	co-dominant	0.89 (0.66-1.20)	0.45
	G/A	94	0.33	112	0.37	dominant	1.02 (0.71-1.45)	0.93
	A/A	17	0.06	9	0.03	recessive	0.35 (0.14-0.85)	0.017

Freq, frequency; OR, odds ratio; CI, confidence interval.

Helix Inc., MT, USA). We assessed the associations between SNPs and SANS scores, using one-way ANOVA to analyze the relationship between SANS scores and genotype frequencies in the schizophrenia subgroup. We calculated the power, given the sample size, using a genetic power calculator <http://pngu.mgh.harvard.edu/~purcell/gpc>[49], and, to reduce error, we adjusted the effective sample size (calculated sample size \times 100/95).

When we calculated the sample powers, we found our case-control study was sufficiently powerful to determine a positive association. In this study, we found sample powers of 0.8046 (rs39315, effective sample size, number of cases for 80% power = 284), 0.8493 (rs17132543, n = 252), 0.9338 (rs3779548, n = 188), 0.9677 (rs733154, n = 155), 0.9145 (rs2024233, n = 208), and 0.9689 (rs4730775, n = 154) for detecting a two-fold increased risk, assuming




Table 2: Primer Sequences

SNP	Sequence(5'-3')	Product Size	Temperature
rs39315	Forward CCTCCCTATGGGCTCTGTATT Reverse CACGGGTGCATGAAATGATGG	450	60
rs17132543	Forward AGCCTCTAGAGAAGTCTGAAG Reverse CTCCTCAACCACTCACACACA	373	60
rs3779548	Forward GTGTGGCCTACTTTGAGAAG Reverse TTCTCCAGCACCTAGACTGTG	355	60
rs733154	Forward GGATCCTTGATCGAGCAGAGCCA Reverse GACTGCAGCAGGAGAGACAGTTA	301	60
rs1051751	Forward TGGGCCACAGAACGAGTATAAC Reverse CCAGAGCTTCCAGGCAGTCTCT	327	65
rs2024233	Forward GTAACAAGTGGGGACGTGTGT Reverse GAGATTCCATGGGTCACATGCA	319	62
rs4730775	Forward TGGGGATACAAGATTGGTGAAC Reverse GATGGCAGAAGCCAACCACTA	360	65
rs6948009	Forward GGTCATTTAGACTGAGACTCG Reverse CACCAATCCCTTCGCCTCTCT	461	60

an α -level of 0.05. However, the sample powers in the divided schizophrenia subgroup (with or without poor concentration) were not sufficient (data not shown). We applied the Bonferroni correction by multiplying P values by the number of SNPs analyzed ($n = 6$).

Results

Of the eight SNPs we examined, all were polymorphic. The genotype distributions of six SNPs (rs39315, rs17132543, rs3779548, rs733154, rs2024233, and rs4730775) were in HWE ($P > 0.05$). We did not estimate one SNP (rs6948009) for the association analysis because its genotype distribution was not in HWE ($p < 0.05$). Also, we excluded one SNP (rs1052751) from the analysis, since it was not polymorphic. Of these SNPs, one SNP (rs4730775) was weakly associated with schizophrenia. Despite this association ($p = 0.017$ in the recessive model), we could not prove the statistical significance of analysis after the Bonferroni correction. Thus, these results do not support a significant role for the *WNT2* sequence variation in the etiology of schizophrenia.




Figure 2 Linkage disequilibrium (LD) blocks of the *WNT2* gene. LD coefficient ($|D'|$) and LD blocks among *WNT2* SNPs. Block 1 consists of rs4730775 and rs2024233. Block 2 comprises rs3779548 and rs17132543.

Table 3: WNT2 SNPs haplotype frequencies in schizophrenia patients and healthy controls.

Haplotype (Block 1)	Type	Schizophrenia		Control			Models	OR(95% CI)	P value	
		Freq	%	Freq	Freq	%				
HAP1 GG	HH	51	0.19		66	0.21		co-dominant	0.92(0.73-1.16)	0.4778
	H-	142	0.52	0.44	163	0.51	0.46	dominant	0.91(0.64-1.31)	0.6236
	--	82	0.3		89	0.28		recessive	0.87(0.58-1.31)	0.5004
HAP2 GA	HH	31	0.11		37	0.12		co-dominant	1.04(0.82-1.32)	0.7506
	H-	121	0.44	0.33	132	0.42	0.32	dominant	1.09(0.79-1.51)	0.604
	--	123	0.45		149	0.47		recessive	0.96(0.58-1.60)	0.8901
HAP3 AA	HH	15	0.05		11	0.03		co-dominant	1.07(0.81-1.42)	0.6325
	H-	93	0.34	0.22	113	0.36	0.21	dominant	1.01(0.73-1.41)	0.9446
	--	167	0.61		194	0.61		recessive	1.61(0.73-3.57)	0.2405

Haplotype (Block 2)	Type	Schizophrenia		Control			Models	OR(95% CI)	P value	
		Freq	%	Freq	Freq	%				
HAP1 AG	HH	50	0.18		65	0.2		co-dominant	0.90(0.72-1.14)	0.3894
	H-	130	0.47	0.42	152	0.48	0.44	dominant	0.88(0.63-1.24)	0.4723
	--	95	0.35		101	0.32		recessive	0.86(0.57-1.30)	0.4881
HAP2 GA	HH	31	0.11		35	0.11		co-dominant	1.01(0.80-1.29)	0.912
	H-	123	0.45	0.34	142	0.45	0.33	dominant	1.01(0.73-1.40)	0.9338
	--	121	0.44		141	0.44		recessive	1.03(0.62-1.72)	0.9181
HAP3 AA	HH	15	0.05		14	0.04		co-dominant	1.12(0.86-1.48)	0.4011
	H-	104	0.38	0.24	114	0.36	0.22	dominant	1.13(0.82-1.57)	0.4568
	--	156	0.57		190	0.6		recessive	1.25(0.59-2.64)	0.5543

Freq, frequency; OR, odds ratio; CI, confidence intervals.

To further analyze the haplotype structure in our samples, we characterized the two LD blocks between the six *WNT2* SNPs (including the 5'UTR and 3'UTR) in the control subjects, using the pairwise *D'* values (Figure 2). Per the criteria used by Gabriel et al. [50], the LD data revealed two haplotype blocks across *WNT2*. The *D'* values from rs4730775 to rs32024233 and from rs37795488 to rs17132543 ranged between 0.95 and 0.99, indicating strong LD between each pair of markers (Figure 2). We performed a haplotype-based association analysis between the schizophrenic and control groups for different combinations of SNPs within *WNT2*, revealing two LD blocks (block 1, rs4730775 and rs32024233; block 2, rs37795488 and rs17132543). However, we did not detect significant associations between the haplotypes in block 1

or 2 and schizophrenia (all *p*-values of the analysis > 0.05; Table 3). We then assessed the gene and haplotype associations with these SNPs in the poor-concentration schizophrenia subgroup (determined using the OPCRIT Checklist). In this subgroup, the genotype distributions of the seven SNPs considered were in HWE (*p* > 0.05). We then analyzed the genetic associations of these SNPs within the subgroup. The SNPs of the *WNT2* gene were not associated with poor concentration, a clinical symptom of schizophrenia (Table 4). In addition, we assessed the associations between SNPs and SANS scores. No statistically significant differences appeared in the total SANS scores among the seven SNPs (Table 5). To compare our genotypic results with different ethnic populations, we searched the human SNP database (<http://>

Table 4: *WNT2* genotype frequencies in Korean schizophrenia patients with poor concentration

SNP	Genotype	NPC		PC		Model	OR (95% CI)	P value
		Freq	%	Freq	%			
rs39315 5'-near gene	A/A	40	33.1	50	29.9	co-dominant	0.97 (0.70-1.34)	0.85
	A/G	52	43.0	85	50.9	dominant	1.16 (0.70-1.92)	0.57
	G/G	29	24.0	32	19.2	recessive	0.75 (0.43-1.33)	0.33
rs17132543 Intron4	A/A	40	33.1	57	34.1	co-dominant	0.99 (0.71-1.38)	0.95
	A/G	59	48.8	79	47.3	dominant	0.96 (0.59-1.58)	0.88
	G/G	22	18.2	31	18.6	recessive	1.02 (0.56-1.87)	0.95
rs3779548 Intron4	A/A	50	41.3	77	46.1	co-dominant	0.90 (0.63-1.27)	0.54
	A/G	57	47.1	71	42.5	dominant	0.82 (0.51-1.32)	0.42
	G/G	14	11.6	19	11.4	recessive	0.99 (0.47-2.06)	0.97
rs733154 Intron4	A/A	67	55.4	104	62.3	co-dominant	0.84 (0.56-1.27)	0.41
	A/G	50	41.3	53	31.7	dominant	0.71 (0.44-1.15)	0.17
	G/G	4	3.3	10	6.0	recessive	1.77 (0.53-5.89)	0.34
rs2024233 3'-UTR	A/A	38	31.4	49	29.3	co-dominant	1.11 (0.79-1.57)	0.53
	A/G	63	52.1	85	50.9	dominant	1.10 (0.66-1.83)	0.72
	G/G	20	16.5	33	19.8	recessive	1.23 (0.67-2.28)	0.50
rs4730775 3'-UTR	G/G	70	57.9	106	63.9	co-dominant	0.87 (0.59-1.28)	0.48
	G/A	45	37.2	49	29.5	dominant	0.76 (0.47-1.24)	0.27
	A/A	6	5.0	11	6.6	recessive	1.29 (0.46-3.62)	0.62
rs6948009 3'-near gene	G/G	87	73.1	124	74.7	co-dominant	0.80 (0.48-1.33)	0.39
	G/A	29	24.4	42	25.3	dominant	0.89 (0.52-1.53)	0.67
	A/A	3	2.5	0	0	recessive	0.00 (0.00-NA)	0.02

Freq, frequency; NPC, nonpoor concentration; PC, poor concentration; OR, odds ratio; CI, confidence interval; N/A, not applicable.

www.ncbi.nlm.nih.gov/SNP dbSNP Build 130). This database presents genotype frequencies for the SNPs analyzed in this study (Table 6). The control group's genotype distributions of the SNPs we analyzed are similar to those of Asian populations, especially the Japanese population (Table 6).

Discussion

The *WNT2* gene and its chromosomal location have received attention as a candidate gene with regard to autism. Given the similarities between autistic disorders and schizophrenia (as a neurodevelopmental disorder), we investigated whether *WNT2* gene variations act as risk factors for schizophrenia in a Korean sample. The results

suggest that *WNT2* may not be involved in the pathogenesis of schizophrenia. Of all the SNPs and haplotypes analyzed, only one SNP (rs4730775) showed a weak association with the disorder. However, the significance disappeared after the Bonferroni correction for multiple testing ($p = 0.102$, Table 1). In addition, we investigated the LD between the *WNT2* SNPs and performed a haplotype analysis between the schizophrenia and control subjects. We could not find any association between *WNT2* and schizophrenia, indicating that there is no genetic association between *WNT2* and schizophrenia.

To further analyze the association test between clinical symptoms and SNP genotypes, we evaluated other symptoms, from the OPCRIT checklist and the total SANS

Table 5: Comparison of SANS scores between schizophrenic patient groups with different genotypes of *WNT2*.

SNP	Genotype	n	SANS		
			Mean	S.E.	P value
rs39315 5'-near gene	A/A	90	66.78	2.627	0.388
	A/G	137	63.76	2.239	0.866
	G/G	61	64.41	2.831	0.55
rs17132543 Intron4	A/A	97	63.99	2.46	0.458
	A/G	138	66.51	2.256	0.278
	G/G	53	62.06	3.016	0.63
rs3779548 Intron4	A/A	127	66.42	2.139	0.689
	A/G	128	65.17	2.257	0.124
	G/G	33	57.48	4.394	0.062
rs733154 Intron4	A/A	171	66.11	1.946	0.371
	A/G	103	63.32	2.375	0.684
	G/G	14	60.5	6.791	0.429
rs2024233 3'-UTR	A/A	87	60.26	2.717	0.031
	A/G	148	67.59	2.034	0.461
	G/G	53	64.68	3.319	0.311
rs4730775 3'-UTR	G/G	176	65.82	1.85	0.267
	G/A	94	64.23	2.63	0.619
	A/A	17	58.82	6.56	0.427
rs6948009 3'-near gene	A/A	211	65.02	1.81	0.981
	A/G	71	64.94	2.5	0.307
	G/G	3	52.33	3.38	0.406

SANS, Scale for the Assessment of Negative Symptoms; S.E., Standard Error; SNP, single nucleotide polymorphism.

scores. However, we found no statistically significant association between any symptom in OPCRIT (Table 4) and total SANS score (Table 5). We compared our genotype frequencies with the human SNP database <http://www.ncbi.nlm.nih.gov/sites/entrez?db=snp>, to show ethnic similarities and differences. The genotype frequencies of our study sample resemble those of the Japanese and Chinese Hapmap populations (Table 6). Further studies are necessary to elucidate (i) whether another case-control study is appropriate and (ii) whether promoter SNPs can affect the expression of *WNT2*. To confirm or refute the lack of association between the *WNT2* gene and schizophrenia, replication studies with adequate sample sizes, or studies with additional SNPs not analyzed in the present study, may be required.

This report has several limitations. First, the sample size may not have been sufficient to detect associations of smaller effects on schizophrenia. We estimated that this sample had more than 0.8 powers to detect association, with the gene exerting a genotypic relative risk of approximately 2. However, schizophrenia appears to be a very genetically complex disease, and the *WNT2* gene risks may be smaller than 2. Second, when we designed the experiment, the versions of the HapMap database and dbSNP database were slightly different compared to their present forms, and, as a result, our selection of SNPs provided incomplete coverage of currently-known common variations in the *WNT2* gene. Furthermore, the present databases showed different locations for some SNP from

Table 6: WNT2 SNPs genotype frequencies in each population

SNP	Genotype	Schizophrenia	Control	Europe	China	Japan	Sub-Saharan African
rs39315 5'-near gene	A/A	0.31	0.26	0.36	0.29	0.46	0.22
	A/G	0.48	0.49	0.41	0.62	0.43	0.44
	G/G	0.21	0.25	0.22	0.09	0.11	0.34
P value				0.285	0.010	0.003	0.373
rs17132543 intron4	A/A	0.34	0.32	0.78	0.24	0.44	0.97
	A/G	0.48	0.48	0.22	0.58	0.47	0.03
	G/G	0.18	0.20	0.00	0.18	0.09	0.00
P value				0.000	0.334	0.048	0.000
rs3779548 Intron4	A/A	0.44	0.44	0.30	0.69	0.36	0.17
	A/G	0.44	0.45	0.47	0.22	0.44	0.65
	G/G	0.11	0.11	0.23	0.09	0.20	0.18
P value				0.031	0.001	0.180	0.000
rs733154 Intron4	A/A	0.59	0.58	No data	0.85	0.62	0.31
	A/G	0.36	0.36	No data	0.13	0.31	0.37
	G/G	0.05	0.06	No data	0.03	0.08	0.33
P value				N/A	0.000	0.675	0.000
rs1051751 Exon 5 Cys294Phe	G/G	1.00	1.00	1.00	1.00	0.98	1.00
	G/T	0.00	0.00	0.00	0.00	0.02	0.00
	T/T	0.00	0.00	0.00	0.00	0.00	0.00
P value				N/A	N/A	N/A	N/A
rs2024233 3'-UTR	A/A	0.30	0.28	0.50	0.29	0.34	0.54
	A/G	0.51	0.51	0.40	0.44	0.50	0.41
	G/G	0.18	0.21	0.10	0.27	0.16	0.05
P value				0.003	0.526	0.531	0.000
rs4730775 3'-near gene	G/G	0.61	0.60	0.37	0.71	0.62	0.46
	G/A	0.33	0.37	0.36	0.27	0.29	0.45
	A/A	0.06	0.03	0.27	0.02	0.09	0.09
P value				0.000	0.261	0.135	0.060
rs6948009 3'-near gene	A/A	0.74	0.53	No data	0.95	0.93	No data
	A/G	0.25	0.45	No data	0.05	0.05	No data
	G/G	0.01	0.02	No data	0.00	0.02	No data
P value				N/A	0.000	0.000	N/A

From database <http://www.ncbi.nlm.nih.gov/sites/entrez?db=snp>. We estimated all p-values via comparisons between the control group of our sample and each population. N/A, not applicable.

previous databases. Consequently, we may have missed some of the relevant associations of *WNT2* genes.

Conclusions

We investigated possible associations between *WNT2* gene SNPs and schizophrenia, and the results were negative. For that reason, it appears the SNPs of *WNT2* may not influence the development of schizophrenia in the Korean population, but, still, additional genetic studies will help develop understanding of the precise mechanisms underlying pathogenesis in schizophrenic patients.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Authors JWK and JHC designed and directed the whole project. Authors SKK and SW K managed the literature searches and analyses. JYS, ARC, and JKP carried out the schizophrenia assessments and advised on patient selection. Author HKP advised on control group collection. Author HJK performed most of the statistical analyses and the genotyping and contributed substantially to the first draft of the manuscript. All authors contributed to and have approved the final manuscript

Acknowledgements

This study was supported by a grant of the Korea Healthcare technology R&D project, Ministry for Health, Welfare & Family Affairs, Republic of Korea (A090239)

Author Details

¹College of Medicine, Soonchunhyang University, Chunan 336-745, Republic of Korea, ²Department of Neuropsychiatry, School of Medicine, Kyung Hee University, Seoul 130-701, Republic of Korea, ³Kohwang Medical Research Institute, School of Medicine, Kyung Hee University, Seoul 130-701, Republic of Korea and ⁴Department of Emergency Medicine, East-West Neo-media Center, Kyung Hee University, 149 Sang-il dong, Gangdong-gu, Seoul 134-727, Republic of Korea

Received: 7 October 2009 Accepted: 24 May 2010

Published: 24 May 2010

References

1. Freedman R: **Schizophrenia.** *N Engl J Med* 2003, **349**:1738-1749.
2. Ross CA, Margolis RL, Reading SA, Pletnikov M, Coyle JT: **Neurobiology of schizophrenia.** *Neuron* 2006, **52**:139-153.
3. Murray RM, Lewis SW: **Is schizophrenia a neurodevelopmental disorder?** *BMJ* 1987, **295**:681-682.
4. Weinberger DR: **Implications of normal brain development for the pathogenesis of schizophrenia.** *Arch Gen Psychiatry* 1987, **44**:660-669.
5. Bloom FE: **Advancing a neurodevelopmental origin for schizophrenia.** *Arch Gen Psychiatry* 1993, **50**:224-227.
6. Lewis DA, Lieberman JA: **Catching up on schizophrenia: Natural history and neurobiology.** *Neuron* 2000, **28**:325-334.
7. Brandon NJ, Millar JK, Korth C, Sive H, Singh KK, Sawa A: **Understanding the role of DISC1 in psychiatric disease and during normal development.** *J Neurosci* 2009, **29**:12768-12775.
8. Freedman R, Goldowitz D: **Studies on the hippocampal formation: From basic development to clinical applications: Studies on schizophrenia.** *Prog Neurobiol* 2009 in press.
9. Potkin SG, Turner JA, Guffanti G, Lakatos A, Fallon JH, Nguyen DD, Mathalon D, Ford J, Lauriello J, Macciardi F, FBIRN: **A genome-wide association study of schizophrenia using brain activation as a quantitative phenotype.** *Schizophr Bull* 2009, **35**:96-108.
10. Nusse R: **Wnt signaling in disease and in development.** *Cell Res* 2005, **15**:28-32.
11. Kozlovsky N, Belmaker RH, Agam G: **GSK-3 and the neurodevelopmental hypothesis of schizophrenia.** *Eur Neuropsychopharmacol* 2002, **12**:13-25.
12. Cotter D, Kerwin R, Al-Sarraj S, Brion JP, Chadwich A, Lovestone S: **Abnormalities of Wnt signaling in schizophrenia - evidence for neurodevelopmental abnormality.** *Neuroreport* 1998, **9**:1379-1383.
13. Miyaoka T, Seno H, Ishino H: **Increased expression of Wnt-1 in schizophrenic brains.** *Schizophr Res* 1999, **38**:1-6.
14. Emamian ES, Hall D, Birnbaum MJ, Karayiorgou M, Gogos JA: **Convergent evidence for impaired AKT1-GSK3beta signaling in schizophrenia.** *Nat Genet* 2004, **36**:131-137.
15. Kozlovsky N, Belmaker RH, Agam G: **Low GSK-3 activity in frontal cortex of schizophrenic patients.** *Schizophr Res* 2001, **52**:101-105.
16. Kozlovsky N, ShanonWeickert C, Tomaskovic-Crook E, Kleinman JE, Belmaker RH, Agam G: **Reduced GSK-3betamRNA levels in postmortem dorsolateral prefrontal cortex of schizophrenic patients.** *J Neural Transm* 2004, **111**:1583-1592.
17. Katsu T, Ujike H, Nakano T, Tanaka Y, Nomura A, Nakata K, Takaki M, Sakai A, Uchida N, Imamura T, Kuroda S: **The human frizzled-3 (FZD3) gene on chromosome 8p21, a receptor gene for Wnt ligands, is associated with the susceptibility to schizophrenia.** *Neurosci Lett* 2003, **353**:53-56.
18. Glatt SJ, Wang RS, Yeh YC, Tsuang MT, Faraone SV: **Five NOTCH4 polymorphisms show weak evidence for association with schizophrenia: evidence from meta-analyses.** *Schizophr Res* 2005, **73**:281-290.
19. Jeong SH, Joo EJ, Ahn YM, Lee KY, Kim YS: **Investigation of genetic association between human Frizzled homolog 3 gene (FZD3) and schizophrenia: Results in a Korean population and evidence from metaanalysis.** *Psychiatry Res* 2006, **143**:1-11.
20. Cotter D, Kerwin R, Al-Sarraj S, Brion JP, Chadwich A, Lovestone S, Anderton B, Everall I: **Abnormalities of Wnt signalling in schizophrenia - evidence for neurodevelopmental abnormality.** *Neuroreport* 1998, **9**:1261-1265.
21. Lovestone S, Killick R, Forti MD, Murray R: **Schizophrenia as a GSK-3 dysregulation disorder.** *Trends Neurosci* 2007, **30**:142-149.
22. Emamian ES, Hall D, Birnbaum MJ, Karayiorgou M, Gogos JA: **Convergent evidence for impaired AKT1-GSK3b signaling in schizophrenia.** *Nat Genet* 2004, **36**:131-137.
23. Miyaoka T, Seno H, Ishino H: **Increased expression of Wnt-1 in schizophrenic brains.** *Schizophr Res* 1999, **38**:1-6.
24. Yang J, Si T, Ling Y, Ruan Y, Han Y, Wang X, Zhang H, Kong Q, Li X, Liu C, Zhang D, Zhou M, Yu Y, Liu S, Shu L, Ma D, Wei J, Zhang D: **Association study of the human FZD3 locus with schizophrenia.** *Biol Psychiatry* 2003, **54**:1298-1301.
25. Zhang Y, Yu X, Yuan Y, Ling Y, Ruan Y, Si T, Lu T, Wu S, Gong X, Zhu Z, et al.: **Positive association of the human frizzled 3 (FZD3) gene haplotype with schizophrenia in Chinese Han population.** *Am J Med Genet B Neuropsychiatr Genet part B* 2004, **129**:16-19.
26. Katsu T, Ujike H, Nakano T, Tanaka Y, Nomura A, Nakata K, Takaki M, Sakai A, Uchida N, Imamura T, Kuroda S: **The human frizzled-3 (FZD3) gene on chromosome 8p21, a receptor gene for Wnt ligands, is associated with the susceptibility to schizophrenia.** *Neurosci Lett* 2003, **353**:53-56.
27. Faraone SV, Matise T, Svrakic D, Pepple J, Malaspina D, Suarez B, Hampe C, Zambuto CT, Schmitt K, Meyer J, Markel P, Lee H, Harkavy Friedman J, Kaufmann C, Cloninger CR, Tsuang MT: **Genome scan of European-American schizophrenia pedigrees: results of the NIMH Genetics Initiative and Millennium Consortium.** *Am J Med Genet* 1998, **81**:290-295.
28. Cadigan KM, Nusse R: **Wnt signaling: a common theme in animal development.** *Genes Dev* 1997, **11**:3286-3305.
29. Uusitalo M, Heikkila M, Vainio S: **Molecular genetic studies of Wnt signaling in the mouse.** *Exp Cell Res* 1999, **253**:336-348.
30. Hauptmann G, Gerster T: **Regulatory gene expression patterns reveal transverse and longitudinal subdivisions of the embryonic zebra fish forebrain.** *Mech Dev* 2000, **91**:105-118.
31. Landesman Y, Sokol SY: **Xwnt-2b is a novel axis-inducing Xenopus Wnt, which is expressed in embryonic brain.** *Mech Dev* 1997, **63**:199-209.
32. Brown JD, Moon RT: **Wnt signaling: why is everything so negative?** *Curr Opin Cell Biol* 1998, **10**:182-187.
33. Wodarz A, Nusse R: **Mechanisms of Wnt signaling in development.** *Ann Rev Cell Dev Biol* 1998, **14**:59-88.
34. Salinas PC: **Wnt factors in axonal remodelling and synaptogenesis.** *Biochem Soc Symp* 1999, **65**:101-109.

35. Wassink TH, Piven J, Vieland VJ, Huang J, Swiderski RE, Pietila J, Braun T, Beck G, Folstein SE, Haines JL, Sheffield VC: **Evidence supporting WNT2 as an autism susceptibility gene.** *Am J Med Genet* 2001, **105**:406-413.
36. Li J, Nguyen L, Gleason C, Lotspeich L, Spiker D, Risch N, Myers RM: **Lack of evidence for an association between WNT2 and RELN polymorphisms and autism.** *Am J Med Genet B Neuropsychiatr Genet* 2004, **126B**:51-57.
37. McCoy PA, Shao Y, Wolpert CM, Donnelly SL, Ashley-Koch A, Abel HL, Ravan SA, Abramson RK, Wright HH, DeLong GR, Cuccaro ML, Gilbert JR, Pericak-Vance MA: **No association between the WNT2 gene and autistic disorder.** *Am J Med Genet* 2002, **114**:106-109.
38. Proitsi P, Li T, Hamilton G, Di Forti M, Collier D, Killick R, Chen R, Sham P, Murray R, Powell J, Lovestone S: **Positional pathway screen of wnt signaling genes in schizophrenia: association with DKK4.** *Biol Psychiatry* 2008, **63**:13-16.
39. American Psychiatric Association: *Diagnostic and statistical manual of mental disorders* 4th edition. Washington DC: American Psychiatric Press; 1994.
40. Flemenbaum A, Zimmermann RL: **Inter- and intra-rater reliability of the brief psychiatric rating scale.** *Psychological Reports* 1973, **32**:783-792.
41. Andreasen NC: **Negative symptoms in schizophrenia. Definition and reliability.** *Archives of General Psychiatry* 1982, **39**:784-788.
42. McGuffin P, Farmer A, Harvey I: **A polydiagnostic application of operational criteria in studies of psychotic illness. Development and reliability of the OPCRIT system.** *Arch Gen Psychiatry* 1991, **48**:764-770.
43. Kim HJ, Park HJ, Jung KH, Ban JY, Ra J, Kim JW, Park JK, Choe BK, Yim SV, Kwon YK, Chung JH: **Association study of polymorphisms between DISC1 and schizophrenia in a Korean population.** *Neurosci Lett* 2008, **430**:60-63.
44. Rickham PP: **Human experimentation. Code of ethics of the world medical association. Declaration of helsinki.** *British Medical Journal* 1964, **2**:177.
45. Solé X, Guinó E, Valls J, Iñiesta R, Moreno V: **SNPStats: a web tool for the analysis of association studies.** *Bioinformatics* 2006, **22**:1928-1929.
46. Barrett JC, Fry B, Maller J, Daly MJ: **Haploview: analysis and visualization of LD and haplotype maps.** *Bioinformatics* 2005, **21**:263-265.
47. Stephens M, Smith NJ, Donnelly P: **A new statistical method for haplotype reconstruction from population data.** *Am J Hum Genet* 2001, **68**:78-89.
48. Jung HY, Park JS, Park YJ, Kim YK, Kimm K, Koh IS: **HapAnalyzer: Minimum Haplotype Analysis System for Association Studies.** *Genomics & Informatics* 2004, **2**:107-109.
49. Purcell S, Cherny SS, Sham PC: **Genetic Power Calculator: design of linkage and association genetic mapping studies of complex traits.** *Bioinformatics* 2003, **19**:149-150.
50. Gabriel SB, Schaffner SF, Nguyen H, Moore JM, Roy J, Blumenstiel B, Higgins J, DeFelice M, Lochner A, Faggart M, Liu-Cordero SN, Rotimi C, Adeyemo A, Cooper R, Ward R, Lander ES, Daly MJ, Altshuler D: **The structure of haplotype blocks in the human genome.** *Science* 2002, **296**:2225-2229.

Pre-publication history

The pre-publication history for this paper can be accessed here:
<http://www.biomedcentral.com/1471-2350/11/78/prepub>

doi: 10.1186/1471-2350-11-78

Cite this article as: Kim *et al.*, No association between polymorphisms of WNT2 and schizophrenia in a Korean population *BMC Medical Genetics* 2010, **11**:78

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

