

CASE REPORT

Open Access

Apparently synonymous substitutions in *FGFR2* affect splicing and result in mild Crouzon syndrome

Aimee L Fenwick^{1†}, Jacqueline AC Goos^{2†}, Julia Rankin³, Helen Lord⁴, Tracy Lester⁴, A Jeannette M Hoogeboom⁵, Ans MW van den Ouweland⁵, Steven A Wall⁶, Irene MJ Mathijssen² and Andrew OM Wilkie^{1,6*}

Abstract

Background: Mutations of *fibroblast growth factor receptor 2 (FGFR2)* account for a higher proportion of genetic cases of craniosynostosis than any other gene, and are associated with a wide spectrum of severity of clinical problems. Many of these mutations are highly recurrent and their associated features well documented. Crouzon syndrome is typically caused by heterozygous missense mutations in the third immunoglobulin domain of *FGFR2*.

Case presentation: Here we describe two families, each segregating a different, previously unreported *FGFR2* mutation of the same nucleotide, c.1083A>G and c.1083A>T, both of which encode an apparently synonymous change at the Pro361 codon. We provide experimental evidence that these mutations affect normal *FGFR2* splicing and document the clinical consequences, which include a mild Crouzon syndrome phenotype and reduced penetrance of craniosynostosis.

Conclusions: These observations add to a growing list of *FGFR2* mutations that affect splicing and provide important clinical information for genetic counselling of families affected by these specific mutations.

Keywords: Craniosynostosis, Crouzon syndrome, Expressivity, *FGFR2*, Penetrance, Splicing, Synonymous substitution

Background

Craniosynostosis defines the premature fusion of the cranial sutures and has an overall prevalence of 1 in 2100–2300 live births [1,2]. Nearly one quarter of craniosynostosis has a genetic aetiology [3,4]; there is considerable genetic heterogeneity and frequent phenotypic overlap between different syndromes. Genes encoding three members of the fibroblast growth factor receptor family (*FGFR1*, *FGFR2* and *FGFR3*) are commonly mutated in individuals with craniosynostosis. Heterozygous mutations in *FGFR2*, which are frequently recurrent, account for ~28% of genetic cases [4] and cause Crouzon [5,6], Pfeiffer [7–9], Apert [10], Beare-Stevenson [11] and bent bone dysplasia [12] syndromes. All involve synostosis of the coronal and other cranial sutures, a distinctive “crouzonoid” craniofacial appearance (comprising hypertelorism,

exorbitism, prominent nose, and midface hypoplasia), but differ in the presence and extent of abnormalities of the hands and feet, other skeletal manifestations and dermatological features [13,14].

FGFR2, like the other members of the *FGFR* family, comprises an extracellular ligand-binding region (composed of three immunoglobulin-like domains), a single transmembrane peptide and a cytoplasmic tyrosine kinase domain. Mutually exclusive alternative splicing of exons IIb and IIc gives rise to epithelial and mesenchymal isoforms (*FGFR2b* and *FGFR2c*) respectively [15]. These alternative extracellular domains interact with different repertoires of fibroblast growth factors (FGFs) to regulate downstream processes such as proliferation, differentiation and cell migration [16].

Here we describe two families heterozygous for the same, previously unreported apparently synonymous variant in *FGFR2* [p.(Pro361Pro)], although caused by differing nucleotide substitutions. The mutation carriers in both families exhibit features of mild Crouzon syndrome, and a minority required craniofacial surgery. We propose that this variant is in fact pathogenic and demonstrate the generation of abnormal cDNA products resulting from

* Correspondence: andrew.wilkie@imm.ox.ac.uk

†Equal contributors

¹Weatherall Institute of Molecular Medicine, John Radcliffe Hospital, University of Oxford, Headington, Oxford OX3 9DS, UK

⁶Craniofacial Unit, Oxford University Hospitals NHS Trust, John Radcliffe Hospital, Oxford, UK

Full list of author information is available at the end of the article

incorrect splicing of exon IIIc in the mutant allele. This finding highlights the challenges posed in interpreting such synonymous variants when providing genetic counselling for affected families.

Case presentations

Family 1

Individual III-1 (Figure 1A) was born after a normal pregnancy and was referred for craniofacial assessment at two years of age because prominent eyes and a head tilt. Her father II-1 (Figure 1B), grandmother I-1 and two deceased uncles were all said to have a similar appearance with prominent staring eyes, but had not required any surgery. She had an occipitofrontal circumference (OFC) of 50 cm (+1.1 SD) and was noted to have a mildly crouzonoid appearance, with slight exorbitism and mid-face retrusion. In view of the mild crouzonoid features, sequencing of *FGFR2* exons IIIa and IIIc was requested. This demonstrated a heterozygous c.1083A>G [p.(Pro361Pro)] variant within exon IIIc, which was also present in II-1 and I-1.

Computed tomography (CT) of the skull of individual III-1 at the age of 2.8 years demonstrated right lambdoid and occipitomastoid synostosis, all other major cranial sutures being patent. Ophthalmological review identified slightly reduced visual acuity and a latent divergent squint with slight left hypophoria. The patient is now four years old and has not undergone any surgical


intervention, as she has a good overall head shape with no major midface retrusion, is making good developmental progress, and has no features to suggest significant intracranial restriction.

Family 2

The male proband (III-2 in Figure 1C), born after an uneventful pregnancy, was referred for craniofacial assessment at the age of three months. Physical examination showed a mild cloverleaf skull with temporal bulging and reduced OFC (36 cm; -2.2 SD), hypertelorism, and severe exorbitism mainly at the infra-orbital level. Skull X-ray and CT showed pansynostosis and multiple craniolacunae, with no intracerebral anomalies.

Owing to the severe peri-orbital features and the absence of deformations of the upper and lower extremities a clinical diagnosis of Crouzon syndrome was suggested. The patient's mother, grandmother and several cousins were reported to show mild facial features also suggestive of this diagnosis.

The proband underwent fronto-orbital advancement at the age of five months. Since the occiput was still severely flattened and both lambdoid sutures were fused, occipital craniotomy and remodelling was performed at the age of twelve months. Clumsiness and motor delay were first noted aged 18 months; psychological testing at the age of 12.8 years gave scores for non-verbal intelligence of 80 (SON-R) and visual-motor integration of 81. Clonidine



was prescribed due to high distractibility and he underwent special education. During childhood, the exorbitism increased requiring further orbital advancement and cranial vault remodelling at the age of eight years. Several deciduous and permanent teeth were extracted because of Class III malocclusion and dental crowding.

Genetic testing of *FGFR2* was performed, identifying the heterozygous point mutation c.1083A>T [p.(Pro361Pro); Figure 2A]. This variant was also present in his mother II-1.

Independently of these events, a second patient (III-1) was referred to the same clinic at the age of ten years with a scaphocephalic head shape. He had previously undergone vault remodelling at the age of 16 months owing to bicoronal synostosis. At the time of referral, he had an occipito-frontal circumference of 48.5 cm (-2.8 SD), hypertelorism and severe exorbitism. In addition, he had

mild maxillary hypoplasia and both 2nd premolars of the lower jaw were absent. Ophthalmic examination showed myopia with divergent strabismus of the right eye associated with reduced visual acuity. A monobloc procedure without distraction (Le Fort III and an advancement of the forehead) was performed.

Due to the severe peri-orbital features a diagnosis of Crouzon syndrome was suggested. Analysis of *FGFR2* identified the heterozygous point mutation c.1083A>T [p.(Pro361Pro)]. His father, grandmother and great-grandmother had a similar craniofacial appearance. Based on the pedigree analysis, it is evident that III-1 and III-2 are third cousins and that the *FGFR2* mutation present in these two branches is identical by descent (Figure 1C). Apart from III-1 and III-2, none of the other affected family members had undergone craniofacial surgery.




Figure 2 Genomic context and consequences of *FGFR2* mutations. **A**, Schematic representation of genomic region affected by c.1083A>G and c.1083A>T mutations (not to scale). IIIa, IIIb and IIIc denote exons of *FGFR2* encoding the 3rd immunoglobulin-like domain; note that physiological skipping of exon IIIb normally occurs in blood mRNA. TM, exon encoding transmembrane domain. Sequencing of genomic DNA demonstrates heterozygosity for c.1083A>G (II-1, Family 1) or c.1083A>T (III-2, Family 2) mutation at the -2 position from the end of exon IIIc (indicated by dashed red line; the upper line in each trace shows the wild type sequence). Below the cartoon is shown the genomic sequence around the cryptic donor splice site within exon IIIc (marked with red asterisk). **B**, Amplified cDNA corresponding to c.1083A>G and c.1083A>T mutations (same individuals as for genomic analyses), demonstrates two additional smaller products (indicated with white arrows **D** and **E**), in addition to the normal (**C**) product. **C**, In the cDNA product coincident with the wild type band (II-1, Family 1), the mutant allele at the penultimate position of exon IIIc is not represented, indicating complete skipping of normal splicing. The consequences for splicing of the mutant allele (sequence traces illustrated are from III-1, Family 2) are either to activate the cryptic splice donor site within exon IIIc (**D**) or to skip exon IIIc completely (**E**).

Materials and methods

Ethics approval for the study was obtained from NRES Committee London - Riverside (09/H0706/20) and the Medical Ethical Committee of the Erasmus University Medical Center Rotterdam (MEC-2013-547). Venous blood was collected into PAXgene Blood RNA tubes (Qiagen) from individuals II-1 (Family 1) and III-1 (Family 2), and RNA was extracted according to the associated protocol. cDNA was synthesized using the Fermentas RevertAid First-Strand Synthesis kit with random hexamer primers according to the manufacturer's instructions.

cDNA was amplified using a forward primer in *FGFR2* exon IIIa (5'-TCGGAGGAGACGTAGAGTTTGTCTG C-3') used in combination with a reverse primer in exon 11 (encoding the transmembrane (TM) domain; 5'-TGT TACCTGTCTCCGCAGGGGATA-3'). DNA bands were cut out and gel purified using the Q-Spin gel extraction kit (Geneflow). Dideoxy sequencing was carried out on the resulting DNA products. The resulting cDNA products were numbered according to NCBI Reference Sequence: NM_000141.4.

Results

The synonymous variants c.1083A>G and c.1083A>T occur at the -2 position of the 5' (donor) splice site of *FGFR2* exon IIIc (Figure 2A). The neural network splice site predictor (http://www.fruitfly.org/seq_tools/splice.html) generates a score for the wild type donor of 0.88, which is reduced to 0.37 by the A>G transition, and to 0.19 by the A>T transversion. In these circumstances, use of a cryptic splice site (score 0.84) within exon IIIc, 51 nucleotides upstream from the end of the exon, is expected based on analysis of a previous mutation c.1084+3A>G [17]. This would lead to an in-frame deletion of 17 amino acids.

Amplification of cDNA from individuals heterozygous for either the *FGFR2* c.1083A>G or the c.1083A>T variants demonstrated the presence of two additional bands, not present in the wild type control, at approximately 430 bp and 330 bp (Figure 2B). Sequencing of the normal 479 bp product from these individuals showed complete

absence of the mutant allele (illustrated in Figure 2C for II-1 from Family 1), indicating that both mutations abolish use of the normal exon IIIc donor splice site. Sequencing of the ~430 bp product confirmed that the cryptic splice donor within exon IIIc was preferred in the mutant allele (Figure 2D), while the ~330 bp product demonstrated complete skipping of exon IIIc (Figure 2E).

Conclusions

In humans there is a consensus sequence around the 5' splice donor site: A, C or T, AG / gtaagt (where / indicates the exon-intron boundary). Mutations of the consensus A at the penultimate nucleotide of the exon have been associated previously with loss of splicing at the donor site [18,19]. Other mutations near the *FGFR2* exon IIIc splice donor site have also been described (Table 1), all associated with a mild Crouzon phenotype. It is likely that the p.(Ala362Ser) substitution [20] in fact exerts its pathogenic effect via the G>T transversion at the -1 position in the exon, rather than as a missense substitution, but this was not tested experimentally.

In the cases reported here, the wild type splice donor site is abolished by the A>G or A>T mutations at the -2 position from the intron, leading to the cryptic site being preferred. The apparently greater amount of mutant cDNA products associated with the A>T mutation appears to correlate both with the greater predicted disruption of the splice site and with the more severe phenotype in clinically affected individuals from Family 2 compared with Family 1. However, since the A>G mutation did not support use of the normal exon IIIc donor splice site (Figure 2C), other explanations for these differences are possible, such as differing proportions of cell types in the blood samples analysed, and/or differences in genetic background.

Utilisation of the cryptic donor would lead to an in-frame deletion of the last 17 amino acids of exon IIIc (p.Gly345_Pro361del), including four residues that form specific contacts with the ligand FGF2 [25]. However as craniosynostosis-causing FGFR mutations function in a

Table 1 Summary of mutations affecting correct splicing of the *FGFR2* exon IIIc donor site

Mutation	Protein	Proposed effect	Experimental demonstration	Reference
c.1032G>A	p.(Ala344Ala)	Activation of cryptic splice site	Yes	Reardon et al. 1994 [6]; Li et al. 1995 [21]; Del Gatto & Breathnach 1995 [22]
c.1083A>G	p.(Pro361Pro)	Loss of normal donor site with use of alternative cryptic splice site	Yes	This study
c.1083A>T	p.(Pro361Pro)	As above	Yes	This study
c.1084G>T	p.(Ala362Ser)	Annotated as missense but likely to affect splicing	No	Everett et al. 1999 [20]
c.1084+3A>G	-	Loss of normal donor site with use of alternative cryptic splice site	Yes	Kan et al. 2004 [17]
c.1084+3A>C	-	Loss of normal donor site with use of alternative cryptic splice site	No	Cornejo-Roldan, Roessler & Muenke 1999 [23]; Kress et al. 2000 [24]

constitutively active dominant manner [26], it is also likely that in these individuals a mutant protein is formed which is prone to forming covalently-linked dimers [26], leading to variable features of Crouzon syndrome.

Our case reports document the range of phenotypic consequences associated with these particular mutations. Whilst a mild crouzonoid phenotype was generally evident, only a minority of individuals developed overt craniosynostosis requiring calvarial surgery. Orthodontic problems may also occur but these were not fully documented in our study. The causes of the clinical variability are unknown, although one potential factor may be the extent of intra-uterine fetal head constraint [27].

In conclusion, mutations near the *FGFR2* exon IIIc splice sites should be carefully evaluated as to whether they may be pathogenic, even if they are synonymous or outside the canonical AG/GT splice acceptor/donor sequences. In particular, the mutations described here are associated with variable Crouzon syndrome features and affected families should be counselled as such.

Consent

Written informed consent was obtained from the patients for publication of this Case report and any accompanying images. A copy of the written consent is available for review by the Editor of this journal.

Competing interests

The authors declare that they have no competing interest.

Authors' contributions

Patient ascertainment and assessment: JACG, JR, AJMH, SAW, IMJM, AOMW. Experimental analysis: ALF, HL, AMWO. Supervised experiments: TL, AOMW. Drafted paper: ALF, JACG, AOMW. All authors approved the paper.

Acknowledgements

We thank all families for their participation, J. Frankland and T. Rostron for DNA sequencing, and W. Baggley for clinical photography. This work was funded by the Wellcome Trust (093329 and 102731) and Oxford NIHR Biomedical Research Centre (A.O.M.W.).

Author details

¹Weatherall Institute of Molecular Medicine, John Radcliffe Hospital, University of Oxford, Headington, Oxford OX3 9DS, UK. ²Department of Plastic and Reconstructive Surgery and Hand Surgery, Erasmus MC, University Medical Center, Rotterdam, The Netherlands. ³Clinical Genetics Department, Royal Devon and Exeter NHS Foundation Trust, Exeter, UK. ⁴Genetics Laboratories, Oxford University Hospitals NHS Trust, Churchill Hospital, Oxford, UK. ⁵Department of Clinical Genetics, Erasmus MC, University Medical Center, Rotterdam, The Netherlands. ⁶Craniofacial Unit, Oxford University Hospitals NHS Trust, John Radcliffe Hospital, Oxford, UK.

Received: 9 June 2014 Accepted: 4 August 2014

Published: 31 August 2014

References

1. Boulet SL, Rasmussen SA, Honein MA: A population-based study of craniosynostosis in metropolitan Atlanta, 1989–2003. *Am J Med Genet A* 2008, **146A**:984–991.
2. Lajeunie E, Le Merrer M, Bonaiti-Pellie C, Marchac D, Renier D: Genetic study of nonsyndromic coronal craniosynostosis. *Am J Med Genet* 1995, **55**:500–504.

3. Wilkie AO, Byren JC, Hurst JA, Jayamohan J, Johnson D, Knight SJ, Lester T, Richards PG, Twigg SR, Wall SA: Prevalence and complications of single-gene and chromosomal disorders in craniosynostosis. *Pediatrics* 2010, **126**:e391–e400.
4. Sharma VP, Fenwick AL, Brockop MS, McGowan SJ, Goos JA, Hoogetboom AJ, Brady AF, Jeelani NO, Lynch SA, Mulliken JB, Murray DJ, Phipps JM, Sweeney E, Tomkins SE, Wilson LC, Bennett S, Cornall RJ, Broxholme J, Kanapin A, 500 Whole-Genome Sequences (WGS500) Consortium, Johnson D, Wall SA, van der Spek PJ, Mathijssen IMJ, Maxson RE, Twigg SRF, Wilkie AOM: Mutations in *TCF12*, encoding a basic helix-loop-helix partner of *TWIST1*, are a frequent cause of coronal craniosynostosis. *Nat Genet* 2013, **45**:304–307.
5. Jabs EW, Li X, Scott AF, Meyers G, Chen W, Eccles M, Mao JI, Charnas LR, Jackson CE, Jaye M: Jackson-Weiss and Crouzon syndromes are allelic with mutations in fibroblast growth factor receptor 2. *Nat Genet* 1994, **8**:275–279.
6. Reardon W, Winter RM, Rutland P, Pulleyn LJ, Jones BM, Malcolm S: Mutations in the fibroblast growth factor receptor 2 gene cause Crouzon syndrome. *Nat Genet* 1994, **8**:98–103.
7. Lajeunie E, Ma HW, Bonaventure J, Munnich A, Le Merrer M, Renier D: *FGFR2* mutations in Pfeiffer syndrome. *Nat Genet* 1995, **9**:108.
8. Rutland P, Pulleyn LJ, Reardon W, Baraitser M, Hayward R, Jones B, Malcolm S, Winter RM, Oldridge M, Slaney SF, Poole MD, Wilkie AOM: Identical mutations in the *FGFR2* gene cause both Pfeiffer and Crouzon syndrome phenotypes. *Nat Genet* 1995, **9**:173–176.
9. Schell U, Hehr A, Feldman GJ, Robin NH, Zackai EH, de Die-Smulders C, Viskochil DH, Stewart JM, Wolff G, Ohashi H, Price RA, Cohen MM Jr, Muenke M: Mutations in *FGFR1* and *FGFR2* cause familial and sporadic Pfeiffer syndrome. *Hum Mol Genet* 1995, **4**:323–328.
10. Wilkie AO, Slaney SF, Oldridge M, Poole MD, Ashworth GJ, Hockley AD, Hayward RD, David DJ, Pulleyn LJ, Rutland P, Malcolm S, Winter RM, Reardon W: Apert syndrome results from localized mutations of *FGFR2* and is allelic with Crouzon syndrome. *Nat Genet* 1995, **9**:165–172.
11. Przylepa KA, Paznekas W, Zhang M, Golabi M, Bias W, Bamshad MJ, Carey JC, Hall BD, Stevenson R, Orlow S, Cohen MM Jr: Fibroblast growth factor receptor 2 mutations in Beare-Stevenson cutis gyrata syndrome. *Nat Genet* 1996, **13**:492–494.
12. Merrill AE, Sarukhanov A, Krejci P, Idoni B, Camacho N, Estrada KD, Lyons KM, Deixler H, Robinson H, Chitayat D, Curry CJ, Lachman RS, Wilcox WR, Krakow D: Bent bone dysplasia-*FGFR2* type, a distinct skeletal disorder, has deficient canonical FGF signaling. *Am J Hum Genet* 2012, **90**:550–557.
13. Agochukwu NB, Solomon BD, Muenke M: Impact of genetics on the diagnosis and clinical management of syndromic craniosynostoses. *Childs Nerv Syst* 2012, **28**:1447–1463.
14. Johnson D, Wilkie AO: Craniosynostosis. *Eur J Hum Genet* 2011, **19**:369–376.
15. Mohammadi M, Olsen SK, Ibrahim OA: Structural basis for fibroblast growth factor receptor activation. *Cytokine Growth Factor Rev* 2005, **16**:107–137.
16. Eswarakumar VP, Lax I, Schlessinger J: Cellular signaling by fibroblast growth factor receptors. *Cytokine Growth Factor Rev* 2005, **16**:139–149.
17. Kan R, Twigg SR, Berg J, Wang L, Jin F, Wilkie AO: Expression analysis of an *FGFR2* IIIc 5' splice site mutation (1084+3A->G). *J Med Genet* 2004, **41**:e108.
18. Podkrajsek KT, Milenkovic T, Odink RJ, der Grinten HL C-v, Bratanic N, Hovnik T, Battelino T: Detection of a complete autoimmune regulator gene deletion and two additional novel mutations in a cohort of patients with atypical phenotypic variants of autoimmune polyglandular syndrome type 1. *Eur J Endocrinol* 2008, **159**:633–639.
19. Twigg SR, Vorgia E, McGowan SJ, Peraki I, Fenwick AL, Sharma VP, Allegra M, Zarakoulis A, Sadighi Akha E, Knight SJ, Lord H, Lester T, Izatt L, Lampe AK, Mohammed SN, Stewart FJ, Verloes A, Wilson LC, Healy C, Sharpe PT, Hammond P, Hughes J, Taylor S, Johnson D, Wall SA, Mavrothalassitis G, Wilkie AOM: Reduced dosage of *ERF* causes complex craniosynostosis in humans and mice and links *ERK1/2* signaling to regulation of osteogenesis. *Nat Genet* 2013, **45**:308–313.
20. Everett ET, Britto DA, Ward RE, Hartsfield JK Jr: A novel *FGFR2* gene mutation in Crouzon syndrome associated with apparent nonpenetrance. *Cleft Palate Craniofac J* 1999, **36**:533–541.
21. Li X, Park WJ, Peyeritz RE, Jabs EW: Effect of splicing of a silent *FGFR2* mutation in Crouzon syndrome. *Nat Genet* 1995, **9**:232.
22. Del Gatto F, Breathnach R: A Crouzon syndrome synonymous mutation activates a 5' splice site within the IIIc exon of the *FGFR2* gene. *Genomics* 1995, **27**(3):558–559.

23. Cornejo-Roldan LR, Roessler E, Muenke M: **Analysis of the mutational spectrum of the FGFR2 gene in Pfeiffer syndrome.** *Hum Genet* 1999, **104**:425–431.
24. Kress W, Collman H, Busse M, Halliger-Keller B, Mueller CR: **Clustering of FGFR2 gene mutations in patients with Pfeiffer and Crouzon syndromes (FGFR2 associated craniosynostosis).** *Cytogenet Cell Genet* 2000, **91**(1–4):134–137.
25. Plotnikov AN, Hubbard SR, Schlessinger J, Mohammadi M: **Crystal structures of two FGF-FGFR complexes reveal the determinants of ligand-receptor specificity.** *Cell* 2000, **101**:413–424.
26. Robertson SC, Meyer AN, Hart KC, Galvin BD, Webster MK, Donoghue DJ: **Activating mutations in the extracellular domain of the fibroblast growth factor receptor 2 function by disruption of the disulfide bond in the third immunoglobulin-like domain.** *Proc Natl Acad Sci U S A* 1998, **95**:4567–4572.
27. Johnson D, Wall SA, Mann S, Wilkie AO: **A novel mutation, Ala315Ser, in FGFR2: a gene-environment interaction leading to craniosynostosis?** *Eur J Hum Genet* 2000, **8**:571–577.

doi:10.1186/s12881-014-0095-4

Cite this article as: Fenwick *et al.*: Apparently synonymous substitutions in *FGFR2* affect splicing and result in mild Crouzon syndrome. *BMC Medical Genetics* 2014 **15**:95.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

